

# DANGEROUS POISON

KEEP OUT OF REACH OF CHILDREN  
READ SAFTY DIRECTIONS BEFORE OPENING OR USING

# LAST SUPPER

## ZINC Phosphide (Broadacre) Mouse Bait

ACTIVE CONSTITUENT: 25g/kg ZINC PHOSPHIDE

For the control of heavy infestations of mice in agricultural situations.

THIS PRODUCT IS NOT TO BE AVAILABLE EXCEPT TO PERSONS AUTHORISED OR LICENSED UNDER THE RELEVANT STATE OR TERRITORY POISONS CONTROL FOR SCHEDULE 7 POISONS.

NET CONTENTS:

100 kg

125 kg

150 kg

160 kg

500 kg

### RISK OF DISEASE, WEED SEEDS AND VOLUNTEER PLANTS FROM UNSTERILISED GRAIN

This product may have been prepared using unsterilized wheat grain that has not been subjected to sterilization by irradiation as indicated below. If prepared using unsterilised grain, the bait product (i) if not taken by mice may germinate and give rise to viable volunteer wheat plants, (ii) may carry fungal diseases and (iii) may contain weed seeds. If the presence of disease, weed seeds or volunteer plants affects crop value it is recommended that fully sterilized bait products are used.

PRODUCT MADE USING UNSTERILISED GRAIN.

UNSTERILISED GRAIN SOURCED FROM: \_\_\_\_\_

#### DIRECTIONS FOR USE

##### RESTRAINTS

DO NOT apply bait to bare ground (including fallow where there is no vegetative cover).

DO NOT apply bait in a trail.

DO NOT apply bait if heavy rain is imminent.

DO NOT apply to the outer 50 m of crop or within 50 m of native vegetation. A 50 m buffer zone in crop must be employed to reduce risk to birds and other non-target animals.

DO NOT apply bait unless monitoring of mouse numbers indicates the potential for crop damage of economic significance.

SITUATION	PEST	RATE	CRITICAL COMMENTS
Grain crops, legume crops, canola, safflower, nut crops and pasture.  (Note: for all situations – only as a post emergent treatment, stubble of harvested crop prior to sowing seed, or when seeds have been sown into crop stubble or other similar ground cover)	Mice	1 kg/ha to achieve an even coverage of 2 – 3 grains/m <sup>2</sup>	Bait is only to be applied by aerial application or accurately calibrated ground application equipment to achieve 2-3 grains/ m <sup>2</sup> . It is imperative that all attempts are made to apply bait as evenly as possible to maximise product efficacy and minimise risk to non-target animals. Last Supper Zinc Phosphide (Broadacre) Mouse Bait will provide a high level of control in most situations. However in extremely high mice densities or areas with significant amounts of alternative feed, sufficient mice may remain to cause damage to crops after treatment. Re-treatment after 14 days may therefore be required in certain circumstances.  It is recommended that pre-treatment estimations of the severity of the mouse problem are performed before applying bait and that post-baiting estimates are conducted to assess the effectiveness of the control operation. For information on the most effective methods of monitoring of mouse populations please contact the government department in your state responsible for vertebrate pest control.  Bait may retain activity for several days after light rain or in damp conditions, and light showers can be tolerated. Maximum performance will however be expected under dry conditions.

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.**

**THIS PRODUCT IS TOO HAZARDOUS FOR USE IN THE HOME GARDEN. DO NOT USE THIS PRODUCT IN THE HOME GARDEN.**

#### WITHOLDING PERIOD

**DO NOT HARVEST ANY CROPS OR ALLOW LIVESTOCK TO GRAZE BAITED AREAS FOR 14 DAYS AFTER APPLICATION.**

#### RISK OF DISEASE, WEED SEEDS AND VOLUNTEER PLANTS FROM UNSTERILISED GRAIN

This product may be prepared using unsterilised wheat grain that has not been subjected to sterilisation by irradiation. The main panel of the label of this product indicates whether this product has been prepared using unsterilised grain. If prepared using unsterilised grain, the bait product (i) if not taken by mice may germinate and give rise to viable volunteer wheat plants, (ii) may carry fungal diseases and (iii) may contain weed seeds. If the presence of disease, weed seeds or volunteer plants affects crop value it is recommended that fully sterilised bait products are used.

#### GENERAL INSTRUCTIONS

Bait should be distributed to minimise exposure to non-target wildlife, especially birds and livestock. It is preferable to place baits late in the day when birds have finished feeding. Particular care should be taken when baits are applied near native vegetation using aircraft as high bait concentrations may occur on the ground beneath aircraft that need to climb steeply. Spilt bait must be collected immediately and applied according to label directions or buried below one metre.

#### PROTECTION OF WILDLIFE, FISH, CRUSTACEANS, THE ENVIRONMENT AND LIVESTOCK

Do not contaminate dams, rivers, streams or waterways with product or the used container.

#### STORAGE AND DISPOSAL

The contents of each container should be used within 3 months of opening. Store in closed original containers, in a dry, cool, well-ventilated area out of direct sunlight. Store in a locked room away from children, animals, food, feedstuffs, seed and fertilisers. Store away from acids, water and any source of heat or ignition.

Dispose of empty containers by crushing and burying below one metre in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots. Excess or unused baits must be buried below one metre. Empty containers and product must not be burnt.

#### SAFETY DIRECTIONS

Poisonous if swallowed. Releases dangerous phosphine gas slowly in moist air and immediately if wet. Do not inhale vapour. Avoid contact with eyes and skin. Keep away from water and liquids. Keep away from naked flames – forms toxic gas. When opening the container and using the product wear elbow-length PVC gloves and a full face respirator with combined dust and gas cartridge or supplied air respirator. Wash hands after use. After each day's use, wash gloves and respirator and if rubber wash with detergent and warm water.

#### FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. Do not give direct mouth to mouth resuscitation if swallowed. To protect rescuer, use air-viva, oxy-viva or one way mask. Resuscitate in a well-ventilated area.

#### SAFETY DATA SHEET

Additional information is listed in the SDS which can be obtained from the supplier.

#### CONDITIONS OF SALE

The use of Last Supper Zinc Phosphide (Broadacre) Mouse Bait being beyond the control of the manufacturer, no warranty expressed or implied is given by Wilhelm Rural Pty Ltd, regarding its suitability, fitness or efficiency for any purpose for which it is used by the buyer, whether in accordance with the directions or not and Wilhelm Rural Pty Ltd accepts no responsibility for any consequence whatsoever from the use of this product.

BN: \_\_\_\_\_

DOM: \_\_\_\_\_

APVMA Approval No 68512/102993B